



Marine Algae of Sert Coasts, Libya

Massoud M. Godeh^a ; Alaa. A. Said^b, Fathalla O. El-Menifi^a and Meftah, M. Zarmouh^c

^a Department of Botany, Faculty of Science, University of Benghazi, Benghazi, Libya

^b Faculty of Science, University of Zagazig, Egypt

^c Faculty of Medicine, University of Musrata, Musrata, Libya

ARTICLE INFO

Article history:

Received 30 January 2017

Revised 19 March 2017

Accepted 23 March 2017

Available online 01 April 2017

Keywords:

Marine algae, R/P, Sert coasts and Mediterranean Sea.

* Corresponding author:

E-mail address: masud.abdgalil@uob.edu.ly

M.M. Godeh

ABSTRACT

The present work was mainly intended to make recent data base of marine algae population at Sert coasts as a first step to evaluate some of their chemical composition, antimicrobial activities and economic importance. Collection, identification and classification of marine algae of Sert coasts at Mediterranean Sea eastern Tripoli were done through some irregular visits during 2008 and 2009. A total of 21 algal species (21 genera) were recorded at the study area. Five species of them (23.81%) were belonging to Chlorophyta (5 families), Six species (28.57%) belonging to Phaeophyta (4 families) and Ten species (47.62%) belonging to Rhodophyta (4 families) with special reference to Cystoseiraceae and Dityotaceae families of Phaeophyta and Corallinaceae and Rhodomelaceae families of Rhodophyta. Most of the recorded algae were safely edible and still needed to know more about their chemical composition and antimicrobial activity.

© 2017 University of Benghazi. All rights reserved.

1. Introduction

Many marine organisms produce chemicals with biomedical properties that are not found, or rare, in terrestrial organisms (Nybakken, 2001) although only 4.5% of Mediterranean Sea flora and fauna are introduced, in total more than 400 species (Boudouresque and Verlaque, 2005). Red and brown algae are mainly used as a human food sources (Rajasulochana, et al., 2009) for several economically useful materials, especially to supplement man diet (Prescott, 1984) and they contain high amounts of carbohydrates, protein, minerals (Rupe'rez and Saura-Calxto, 2001) low fat contents, few calories (Lahaye and Kae, 1997) and some of them are considered great potential food sources and present almost year round (Abdallah, 2007). Many metabolites isolated from marine algae have been shown to possess bioactive efforts (Oh et al., 2008, Venkateswarlu et al.,

2007 and Rui-yun Yang et al., 2006). Taxonomy is the hidden but important foundation when seaweeds would be exploited for any of their natural products, so in his work he collected the marine algae of Sert coasts to evaluate the chemical, economic, pharmaceutical importance in further studies.

2. Materials and methods

2.1 The Study area:

Sert coasts and its geographical location and main morphometric characteristics were illustrated at Fig. 1. It lies at 32° 03' 28.03" N and 20° 02' 29.18" E at the western Libyan coast. It characterized by long, curved and mainly sandy coasts with some rocky parts at Mediterranean Sea. It also had commercial, fishing ports and many public beaches.



Fig. 1. Maps of Libya and the study area.

2.2 Sampling and sample preparations:

For the long distance between the study area and research laboratories of Benghazi, specimens were collected through some irregular visits during 2008 and 2009 in ice tanks at polyethylene bags. Some of them sprinkled with 4% formalin sea water solution for mounting on the herbarium sheets, glass bottles and some of them were kept freshly at ice or refrigerators for future use and subsequent taxonomic identification using [Nizamuddin et al., \(1979\)](#), [Burrows \(1991\)](#), [Godeh, et al., \(1992\)](#) and [Aleem \(1993\)](#). Epiphytes, impurities and salts are carefully and quickly removed at laboratory with tap and distilled waters. The herbarium sheets have been deposited in the Herbarium of Botany Department, Garyounis University, Benghazi {CHUG nos. FM. 650; 651}. Longitudinal and transverse sections of the axis at the apexes, midfronds and the bases were handmade and stained in 1% KI₂ or aniline blue solution.

2.3 Species richness:

Species richness index is calculated according to [Wilhm, \(1975\)](#) just by direct count of different green algal species (taxa) at every sampling site where, the decrease in number of species and increase in number of individuals are a characteristic feature of polluted water.

3. Results and discussion

Regarding to species richness, a total of 21 marine green algal species (21 genera) were recorded from some sampling stations along the coasts of Sert city to cover the study area ([Table 1](#)). Five species of them (23.81%) were belonging to Chlorophyta (5 families), Six species (28.57%) belonging to Phaeophyta (4 families) and Nine species (47.62%) belonging to Rhodophyta (4 families).

Regarding to Chlorophyta (5 families), Sert coasts were similar only to Tolmeta coasts which also had 6 green algal species ([Said, et al., 2010](#)). It is relatively poor of green algae as compared to all eastern coasts like Tobruk and Ain-Ghazala coasts which had 14 and 9 species, respectively ([Godeh et al., 2009](#)). Derna and Susa also had 7 and 18 green algal species, respectively ([Said, et al., 2010](#)). Tukra coast had 11 green algal ([Said and Godeh, 2008](#)). Some of the recorded green algae were recorded also at the Turkish Urla coast ([Tüney et al., 2006](#)). At invaded Bay of Marseilles (France, NW Mediterranean) [Klein and Verlaque, \(2007\)](#) identified only 38 green algal taxa belonging to Chlorophyta. Rhodes Island had 33 green macroalgal taxa ([Tsiamis, et al., 2007](#)).

The relatively reduction of green species may be due to the presence of some Caulerpales which considered strong competitors ([David et al., 2004](#) and [Piazzi et al., 2005](#)) and its production of toxic substances, which inhibit their grazing ([Piazzi et al., 2005](#)). The storms also act as an important structuring force on the seagrasses system and sedimentation ([Schlacher-Hoenlinger and Schlacher 1998](#)). [Zimmermann, \(2006\)](#) evaluated the depth, turbidity and light intensity as major limiting factors to the seagrasses growth and development.

[Table 1](#) illustrated 6 identified brown algal species (6 genera) of the present study area of Sert coasts with dominance of Cystoseiraceae and Dictyotaceae families. Some of the Seagrass have been described as excellent bioindicators ([Pergent-Martini et al., 2005](#)) like *Cystoseira* which considered very good indicators of good environment ([Arévalo, et al., 2007](#)) due to their high sensitivity to changes of environment and specifically to human impacts ([Short and Wylline-Echeverria, 1996](#)).

Generally, Sert coasts were also cheaper in Phaeophyceae than the eastern coasts of Libya like Derna and Tolmeta which had 14 brown algal species of each ([Said, et al., 2010](#)) and Tobruk coasts which had 11 and brown algal species ([Godeh, et al., 2009](#)). The study area were more or less similar to Tukra coasts which had 7 brown algal species ([Said, and Godeh, 2008](#)) while it was richer

than Ain-Ghazala coasts which had just one brown species ([Godeh, et al., 2009](#)).

Table 1: List of marine algae at Sert coasts.

Division	Family	Algae
Chlorophyta (5 Families & 5 species)	Anayamaceae	<i>Anadyomene</i> Lamouroux 1812
		<i>Anadyomene stellata</i> (Wulfen) C. Agardh
	Caulerpaceae	<i>Caulerpa</i> Lamouroux 1809
		<i>Caulerpa prolifera</i> (Forsskål) Lamouroux
	Ulvaceae	<i>Enteromorpha</i> Link 1820
		<i>Enteromorpha intestinalis</i> (Linnaeus) Greville
	Udoteaceae	<i>Halimeda</i> Lamouroux 1816
		<i>Halimeda tuna</i> (Ellis et Solander) Lamouroux
	Microdictyaceae	<i>Microdictyon</i> Decaisne 1839
		<i>Microdictyon tenuius</i> (C. Agardh) Decaisne
Phaeophyta (4 Families & 6 species)	Cystereiraceae	<i>Cystoseira</i> C. Agardh 1820
		<i>Cystoseira discors</i> (Linn.) C. Agardh emend Sauvageau
		<i>Padina</i> Adanson 1763
		<i>Padina pavonica</i> (Linnaeus) Lamouroux
	Dictyotaceae	<i>Dictyopteris</i> Lamouroux 1809
		<i>Dictyopteris membranacea</i> (Skackhouse) Batters
		<i>Dictyota</i> Lamouroux 1809
		<i>Dictyota dichotoma</i> (Hudson) Lamouroux
	Stypaulaceae	<i>Halopteris</i> Kützting 1843
		<i>Halopteris scoparia</i> (Linnaeus) Sauvageau
	Sargassaceae	<i>Sargassum</i> C. Agardh 1820
		<i>Sargassum acinarium</i> C. Agardh
Rhodophyta (4 Families & 10 species)	Corallinaceae	<i>Fosliella</i> Howe 1920
		<i>Fosliella zonalis</i> (Crouan frat) Feldmann
		<i>Lithophyllum</i> Philippi
		<i>Lithophyllum solutum</i> (Foslie) Lemoine
		<i>Neogoniolithon</i> Setchell et Mason 1943
		<i>Neogoniolithon mamiulusum</i> (Hauck) Hewe
		<i>Pseudolithophyllum</i> Lemoine 1913
		<i>Pseudolithophyllum expansum</i> (Philippi) Lemoine
	Rhodomelaceae	<i>Laurencia</i> Lamouroux 1813
		<i>Laurencia paniculata</i> (C. Tgaro) Greville
		<i>Lophocladia</i> Schmitz 1896
		<i>Lophocladia lallemandii</i> (Montagne) Schmitz
		<i>Rytiphlaea</i> C. Agardh 1824
		<i>Rytiphlaea tinctoria</i> (Clemente) C. Agardh
		<i>Vidalia</i> Lamouroux 1822
	Peyssonneliaceae	<i>Vidalia volubilis</i> (Linn.) J. Agardh
		<i>Peyssonnelia</i> Decaisne 1841
	Helmethoclaiceae	<i>Peyssonnelia rubra</i> (Greville) J. Agardh
		<i>Liagora</i> Lamouroux 1812
		<i>Liagora viscida</i> (Forsskål) C. Agardh
Total	13 families	21 algal species (21 algal genera)

A total of nine species marine red algal species (47.62%) were recorded at the present study area of Sert coasts and tabulated in [Table 1](#) with special reference to Corallinaceae and Rhodomelaceae families. The recorded red algal species were quite differ than the western Libyan coasts like Tukra coast (38 species) and Tolmeta coasts (13 species) recorded by [Said and Godeh, \(2008\)](#) and [Said, et al., \(2010\)](#), respectively. But more than those recorded at Derna and Susa (just 5 and 6 species,

respectively) by Said, et al., (2010) and those of Tobruk coast (16 species) which recorded by Godeh, et al., (2009).

The result was quite quantitatively differing to South Aegean Sea (Greece) which qualitatively dominated by 60 red algal taxa (Diapoulis and Tsiamis, 2007). Klein and Verlaque, (2007) identified 125 Rhodophytes at invaded Bay of Marseilles (France, NW Mediterranean) and Tsiamis et al., (2007) were studied 95 Rhodophytes on rocky shores of Rhodes Island (Greece). Cinelli, et al., (2007) recorded 62 species or Rhodophyta along the Tuscany coast in north-western Mediterranean Sea.

R/P ratio of Sert coasts was 1.67 which less than the mean value of Tunisian coasts (2.99) and all Mediterranean coasts which ranged between 2.0 and 3.8 (Ben Maiz, et al., 1987).

Although some good indicators algae were recorded, one could conclude that the study area is relatively poor of marine algae and polluted according to Wilhm, (1975) who reported that, the decrease in number of species and increase in number of individuals is a characteristic feature of polluted water. This may be also due to the sandy nature of Sert coasts and differences of sampling programs. Many other factors like continuous distribution of invasive species from many places of the world also affected (Klein and Verlaque, 2007 and many recent reports).

Acknowledgement

Deep thank for the research and consultancies center of Garyounis University for wonderful technical and financial support and all their kind and continuous assistance.

References

- Abdallah, M. A. M., (2007): Chemical composition and trace element concentration of marine algae *Enteromorpha compressa* as a potential food source. Proceeding of the 3rd Mediterranean symposium on marine vegetation. 27-29 March 2007 - Marseilles. 225-228.
- Aleem, A. A. (1993) The marine algae of the Alexandria, Egypt.
- Arévalo, R.; Pinedo, O. and Ballesteros, E. (2007): Changes in the composition and structure of Mediterranean rocky-shore communities following a gradient of nutrient enrichment: descriptive study and test of proposed methods to assess water quality regarding macroalgae. *Marine Pollution Bulletin*, 55: 104-113.
- Ben Maiz, N.; Boudouresque, C. F. and Ouahchi, F. (1987) Inventaire des algues et phanerogames marines benthiques de la Tunisie. Istratto da "Giornale Botanico Italiano" Vol. 121, n. 5-6, 259-304.
- Boudouresque, C. F. and Verlaque, M. (2002) Assessing scale and impact of ship-transported alien macrophytes in the Mediterranean Sea. In: CIESM (ed) "Alien Marine Organisms Introduced by Ships in the Mediterranean and Black Seas" Monaco. CIESM Workshop Monographs, 20: 53-61.
- Burrows, E. M. (1991) Seaweeds of the British Isles. Volume 2. Chlorophyta. London: British Museum (Natural History).
- Cinelli, F., Balata, D. and Piazzzi, L., (2007) Threats to coralligenous assemblages: sedimentation and biological invasions. Proceeding of the 3rd Mediterranean symposium on marine vegetation. Marseilles. 27-29 March 2007 - 42-47.
- David, B.; Luigi, P. and Francesco, C. (2004) A Comparison Among Assemblages in Areas Invaded by *Caulerpa taxifolia* and *C. racemosa* on a Subtidal Mediterranean Rocky Bottom. *Marine Ecology*, 25 (1): 1-13.
- Diapoulis, A. and Tsiamis, K. (2007) Marine flora and vegetation of South Aegean Sea (Greece). Proceeding of the 3rd Mediterranean symposium on marine vegetation. Marseilles. 27-29 March 2007 - 263-264.
- Godeh, M.; El-Menifi F. and Said, A. (2009) Marine algae of Tobruk and Ain Ghazala coasts, Libya. Garyounis University Press.
- Journal of Science and Its Applications. Vol. 3, No. 1, pp 42-55, April 2009.
- Godeh, M.; Nizamuddin, M. and El-Menifi F. (1992) Marine algae from eastern coast of Libya (Cyrenaica). *Pak. J. Bot.*, 24(1): 11-21.
- Ki-Bong Oh, Ji Hye Lee, Soon-Chun Chung, Jongheon Shin, Hee Jae Shin, Hye-Kyeong Kim, Hyi-Seung Lee, (2008) Antimicrobial activities of the bromophenols from the red alga *Odonthalia corymbifera* and some synthetic derivatives, *Bioorganic & Medicinal Chemistry Letters*, 18, 104-108.
- Klein, J. and Verlaque, M. (2007) *Caulerpa racemosa* effect on macrophyte assemblages of dead *Posidonia* beds. Proceeding of the 3rd Mediterranean symposium on marine vegetation. 27-29 March 2007 - Marseilles. 78-82.
- Lahaye, M. and Kae A. B., (1997) Seaweed dietary bres: Structure, physico-chemical and biological properties relevant to intestinal physiology. *Science des Aliments*, 17: 619-639.
- Nizamuddin, M.; West, J. A. and Menez, E. G. (1979): A list of marine algae from Libya. *Bot. Mar.*, 22: 465-476.
- Nybakken, J. W. (2001) *Marine Biology*. An ecological approach. Fifth edition. Benjamin Cumings, an imprint of Addison Wesley Longman, Inc. San Francisco. Pp. 236-308.
- Pergent-Martini, C.; Leoni, V.; Pasqualini, V.; Ardizzone, G. D.; Bedini, R.; Belluscio, A.; Belsher, T.; Borg, J.; Boudouresque, C. F.; Boumaza, S.; Bouquegneau, J. M.; Bula, M. C.; Calvo, S.; Cebrian, J.; Charbonnel, E.; Cinelli, F.; Cossu, A.; Di Maida, G.; Dural, B.; Francour, P.; Gobert, S.; Meinesz, A.; Molenaar, H.; Mansour, H. M.; Panayotidis, P.; Pergent, G.; Piazzzi, L.; Pirrotta, M.; Relini, G.; Romero, J.; Sanchez-lizaso, J. L.; Semroud, R.; Shembri, P.; Shili, A.; Tomasello, A. and Velimirov, B. (2005) Descriptors of *Posidonia oceanica* meadows: Use and Application. *Ecological Indicators*, 5: 213-230.
- Piazzzi, L.; Meinez, A.; Verlaque, M.; Akc.Ali, B.; Antolic, B.; Argyrou, M.; Balata, D.; Ballesteros, E.; Calvo, S.; Cinelli, F.; Cirik, S.; Cossu, A.; D'Archino, R.; Djellouli, S. A.; Javel, F.; Lanfranco, E.; Mifsud, C.; Pala, D.; Panayotidis, P.; Peirano, A.; Pergent, G.; Petrocelli, A.; Ruitton, S.; Zuljvic, A. and Ceccherelli, G. (2005) Invasion of *Caulerpa racemosa* var. *cylindracea* (Caulerpa, Chlorophyta) in the Mediterranean Sea: an assessment of the spread. *Cryptogam. Algol.*, 26: 189-202.
- Prescott, G. W., (1984) Economic of algae. In: "The algae: A Review". Otto Koeltz Science publishers D-629 Koenig-Stein, W. Germany.
- Rupe'rez, P. and Saura-Calxto, F. (2001) Dietary fibre and physicochemical properties of edible Spanish seaweeds. *Europe. J. of Food Res. and Tech.*, 212:349-354.
- Said, A. and Godeh, M. (2008) Marine algae of Tukra and Tolmeta coasts, Libya. *Egyptian J. of Phycol.* Vol. 9, 2008. 167-179.
- Said, A., Godeh, M. and El-Menifi, F. (2010) Marine algae of Derna, Susa and Tolmeta coasts, Libya. The Second International Conference on Phycology, Limnology and Aquatic Sciences. 14-15 February (2010) Port Said, Egypt.
- Schlacher-Hoenlinger, M. A. and Schlacher, T. A. (1998) Accumulation, contamination and seasonal variability of trace metals in the coastal zone-patterns in a seagrass from the Mediterranean. *Mar. Biol.*, 131:401-410.
- Short, F. T. and Wylline-Echeverria, S. (1996) Natural and human-induced disturbances of seagrasses. *Environ. Conserv.*, 23 (1): 17-27.
- Tsiamis, K.; Panayotidis, P. and Montesanto, B. (2007) Contribution to the study of the marine vegetation of Rhodes Island (Greece). Proceeding of the 3rd Mediterranean symposium on marine vegetation. 27-29 March 2007 - Marseilles. 190-196.

- Tüney, I.; Çadırclı, B. H.; Ünal, D. and Sukatar, A. (2006) Antimicrobial Activity of the Extracts of Marine Algae from Coast of Urla (Izmir, Turkey). *Turk. J. Biol.* 30: 171-175.
- Wilhm, J. L. (1975) Biological indicators of pollution- In: Whitton, B. A. (ed.), *River ecology*. - Blackwell. Oxford: pp. 375-400.
- Venkateswarlu, S. Panchagnula, G. K.; Gottumukkala, A. L. and Subbaraju, G. V. (2007) Synthesis, structural revision, and biological activities of 4'-chloroaurone, a metabolite of marine brown alga *Spatoglossum variable*, *Tetrahedron*, 63, 29, 6909-6914.
- Rui-yun Yang, Chun-yuan Li, Yong-cheng Lin, Guang-tian Peng, Zhi-gang She, Shi-ning Zhou, (2006) Lactones from a brown alga endophytic fungus (No. ZZF36) from the South China Sea and their antimicrobial activities, *Bioorganic & Medicinal Chemistry Letters*, 16, 16, 4205-4208.
- Zimmermann, R. C. (2006) Light and photosynthesis in seagrass meadows. In: Larkum W. D. et al., (eds). *"Seagrasses: biology, ecology and conservation"*, Springer, Netherlands: 303-321.