



Faculty of Science - University of Benghazi

Libyan Journal of Science & Technology

journal home page: www.sc.uob.edu.ly/pages/page/77

A conifer *Pagiophyllum yafraiensis* sp. nov. from the Jurassic Khashm az Zarzur Formation, Yafran, Jebel Nefusa – Libya.

Ahmed M. Muftah*, Ashor Y. El Zouki

University of Benghazi, Faculty of Science, Department of Earth Sciences, Benghazi-Libya.

Highlights

- *Pagiophyllum yafraiensis* sp. nov. from the Jurassic Khashm az Zarzur Formation was introduced.
- The age of Khashm az Zarzur Formation was assigned to (Callovian-Oxfordian) in age
- A warm humid environment was interpreted due to the presence of *Pagiophyllum* and *Brachyphyllum* conifers flora, this is also supported sedimentologically.

ARTICLE INFO

Article history:

Received 07 November 2020

Revised 19 October 2021

Accepted 20 October 2021

Keywords:

Pagiophyllum, Jurassic, Fossil conifer, Khashm az Zarzur, Jebel Nefusa, western Libya

*Address of correspondence:

E-mail address: ahmed.alkowafi@uob.edu.ly

A.M. Muftah

ABSTRACT

The Jurassic flora from the Khashm az Zarzur Formation in the vicinity of Yafran village in Jebel Nefusa, northwestern Libya was studied. Among which a new species of fossil conifer, *Pagiophyllum yafraiensis* sp. nov. was described and illustrated.

This new species occurred in association with *Piazopteris branerii*, *Brachyphyllum* sp., *Hirmerella* sp., *Otozamites* sp., and *Samaropsis* sp. According to the stratigraphic position of the Khashm az Zarzur Formation dated as (Callovian-Oxfordian), however, a Late Jurassic (Bathonian-Tithonian) age has been assigned palynologically.

The global paleogeographical distribution of the genus *Pagiophyllum* is also illustrated herein.

1. Introduction

Jebel Nefusa is one of the major geological features in north-western Libya. It is a prominent escarpment rising sharply from a flat plain (the Jefarah) about 75 km, south of Tripoli, as an east-west ridge separating the Ghadames Basin from the Jifarah Basin (El Zouki, 1980a) (Fig. 1). This uplift extends for 400 km from Misratah to the Tunisian border and continues in Tunisia (Hallett, 2002). Fieldwork has been undertaken during the year 1974. Plant fossils were collected from the upper part of Khashm az Zarzur Formation "which was named Chameau Mort Formation" in two sections, the first section (S10) just below Eghaish village and the other section in a sand quarry (S11) just below Yafran town in Al Jebel Nefusa, Northwestern Libya (Fig. 1). These two sections provide the basis upon which the collected fossil plant of this formation has been studied. Rare studies are published in the fossil plants of the Libyan Jurassic-Cretaceous world.

Photographs and latex replicas were prepared in the British Museum. The illustrated specimens including the *Pagiophyllum yafraiensis* sp. nov. are deposited at the paleontology cabinet of the Geological Museum of the Earth Sciences Department, University of Benghazi, Benghazi- Libya.

2. General Stratigraphy

The Mesozoic rocks exposed in the Jebel Nefusa range in age from Late Triassic to Late Cretaceous (Figs. 2, 3). The stratigraphy

of the area is complicated as different nomenclatures are applied by different authors as in Fig. 3, this problem needs more stratigraphical investigations to unify the rock names of the exposed rock units in the region. The Jurassic rocks exposed in the area are arranged into two groups according to El Zouki (1980a), which are from bottom to top (Fig. 2):

- Bir el Ghenem Group (Lower to Middle Jurassic), comprising the Bu Gheilan Limestone - Bir el Ghenem gypsum Formation; Bu en Niran Formation; and Abreghe Formation.
- Tigi Group (Middle to Upper Jurassic), comprising of the Tacbal Limestone and Khashm az Zarzur Formation of El Hinnawy and Cheshitev (1975) "previously Giosc Shale and Chameau Mort Sandstone" and Shakshuk Formation. On the other hand, the Lower Cretaceous is represented by the Cabao (Kabaw) Formation of the Weldean age and the Albian Chicla (Kiklah) Formation.

The Upper Cretaceous is represented by:

- Nefusa Group (Cenomanian), comprising the Ain Tobi Formation, Jefern Marl Formation, and Garian Formation. These three formations of this group are Cenomanian in age.
- Tigrinna Formation (Cenomanian-Turonian) is the youngest Formation in the area.




Fig. 1. Location map of the Jebel Nefusa area, northwestern Libya, the two plant fossil collection sites are indicated by Red dots (modified after El Zouki, 1980a).




Fig. 2. A stratigraphic relationships of the Mesozoic rocks, Jebel Nefusa area (Modified after Cate and Hammuda, 1969; El Zouki, 1976, 1980a; Hallett 2002). The index map after Shiref and Salaj (2007)

2.1 Stratigraphy of the Upper Jurassic and Lower Cretaceous

The upper Jurassic in the area is represented by the Khashm az Zarzur Formation and the Shakshuk Formation (Figs. 2, 3). The Khashm az Zarzur Formation "formerly Chameau Mort Formation of Burolet (1963)" was named by El Hinnawy and Cheshitev for the two rock units overlying the Tekbal Formation which were formerly known as the Giosc Shale and the Chameau Mort Sandstone (Banerjee, 1980). The Khashm az Zarzur Formation is exposed in the scarp of the Jabal Nafusa from the vicinity of Kiklah village in the east to Wazin village in the West at the Tunisian border. The lower unit is lacustrine claystone with traces of salt and gypsum,

with freshwater pelecypods, plant remains, and fossil wood. The upper unit, however, is a continental sandstone as evidenced by SEM analysis of the sand grains with marked cross beddings and occasional claystone bands that deposited close to the shoreline containing plants fossils (El Zouki, 1980b; Hallett, 2002), it comprises multicolored and gypsiferous shale and mudstone with ferruginous organic-rich laminae, thin beds of marl and containing abundant plant fossils. In Yafran and Kiklah areas the Khashm az Zarzur Formation is overlain by the Kiklah Formation and underlain by the Tacbal Formation. In Jado area, the base of the Khashm az Zarzur Formation is unexposed but attains thickness over 30m and it is overlain by the Shakshuk Formation.

Age	Region	Al Hamadah al Hamra (After Shiref & Salaj, 2007)		Hon Graben and Gebel Nafusa (After Barr & Weegar, 1972)		Northwest Libya	
		(Hallett, 2002)		(Fatmi et al, 1978) *			
Cretaceous	Maastrichtian	Al Gharbiyah Fm.		Lower Tar Marl		Al Gharbiyah Fm.	
	Campanian	Mizdah Fm. Thala Mbr.		Thala Mbr.		Mizdah Fm. Thala Mbr.	
	Santonian	Mazuzah Mbr.		Mazuzah Ls. Mbr.		Mazuzah Mbr.	
	Coniacian	Qasr Tigrinnah Marl Fm.		Tigrinna Marl Mbr.		Qasr Tigrinnah Fm.	
	Turonian	Nalut Fm.		Nalut Fm.		Nalut Fm.	
	Cenomanian	Sidi As Sid Fm. Yefren Marl Mbr. Ain Tobl Mbr.		Nefusa Group Gharah Ls. Jefren Marl Ain Tobl Ls.		Sidi as Sid Fm. Yefren Fm. Ain Tobl Fm.	
	Albian	Kiklah Fm.		Kiklah Fm.		Kiklah Fm.	
	Aptian						
	Barremian						
	Hauterivian						
Jurassic	Valanginian						
	Berriasian						
	Tithonian						
	Kimmeridgian						
	Oxfordian	Ar Rajban Fm.					
	Callovian	Shakshuk Fm.					
	Bathonian	Khashm Az Zarzur Fm. Takbal Fm.					
	Bajocian	Abreghs Fm.					
	Aalenian						
	Toarcian	Bir Al Ghanam Bu en Niran Mbr.					
Triassic	Plensbachian						
	Sinemurian						
	Hettangian						
	Rhaetian						

Fig. 3. Different Nomenclatures used by different authors for the Jurassic- Cretaceous deposits in the NW region of Libya (* IRC = Mann 1975; Smetana 1975; El-Hinnawy and Cheshitev 1975; Antonovic 1977; Novović 1977); Burolet 1977; Fatmi 1977; El Zouki 1976; Hammuda 1969; Desio et al 1963; Fatmi et al., 1980; and Christie 1955).

3. Systematic Paleontology

Kingdom: Plantae Haeckel, 1866

Phylum: Pinophyta Reveal, 1996

Class: Pinopsida Burnett, 1833

Order: Pinales Dumortier, 1829

Family: Araucariaceae Henckel and Hochstetter, 1865

Genus: *Pagiophyllum* Heer, 1881

Pagiophyllum (Pomel) Saporta, 1873

Definition: This fossil conifer foliage has been variously assigned to several different conifer groups including Araucariaceae and Cheirolepidiaceae Taylor et al. (2009). Most *Pagiophyllum* species have affinities with other conifers such as the Cheirolepidiaceae and Taxodiaceae (Harris, 1979). *Pagiophyllum* leaves are helically arranged with length exceeds width (Stockey, 1982). They were found around the globe during the Permian to the Cretaceous period <https://en.wikipedia.org/wiki/Pagiophyllum>. The geographical distribution of the genus during the Jurassic world is shown in Fig. 6 (e.g. South Mexico, Cuba, Brazil, Columbia, North Yorkshire of England, North America, and Australia).

Pagiophyllum yafraiensis sp. nov.

(Figs. 4, 5 a-c)

Diagnostic features of the species: It is diagnosis by having papillae on the ordinary epidermal cells of the leaf, quite unlike the other known species.

Etymology of the genus: After Yafran city, where the specimens are found at the Jebel Nefusa escarpment northwest of Libya,

Description: *Pagiophyllum yafraiensis* sp. nov. is a well-preserved conifer shoot of about 2 cm long. The leaves are spirally arranged and variable in shape but they are always more than twice as long as broad (Fig. 5a-c). The scanning electron microscope has shown that the surface of the leaf is covered with tiny papillae and well-defined rows of stomata. The stomata of this new species surrounded by papillose subsidiary cells are very similar to those of the *Pagiophyllum kurrii* Schimper and *P. connivens* Kandall. In general, the dense covering of papillae often indicates that the plant grew in a dry habitat.



Fig. 4. *Pagiophyllum yafraiensis* sp. nov. from Khashm az Zarzur Formation, Jebel Nefusa, Yafran in Libya.




Fig. 5. SEM photos of *Pagiophyllum yafraiensis* sp. nov. from Khashm az Zarzur Formation, Jebel Nefusa, Libya.

Occurrence: The conifer *Pagiophyllum* is recorded from all the continents of the old Jurassic land (Pangea) in the forms of *P. burmense*, *P. marwarensis*, *P. peregrinum*, *P. rewaensis*, *P. maculosum*, *P. connivens*, *P. compactum*, and *P. spinosum* (Fig. 6). However, In this study, the newly introduced species *Pagiophyllum yafraiensis* sp. nov. reported from Jurassic age in the Khashm az Zarzur Formation in the vicinity of Yafran city, Jebel Nefusa the northwestern region of Libya.

Associated assemblage: The *Brachyphyllum* sp. (Figs. 7a, b) A conifer leafy twig of about 1cm long. The spirally arranged leaves here are about twice as long as broad, shorter than in *Pagiophyllum*. The leaves of this species are also densely covered by well-defined rows of stomata (Fig. 7b). The stratigraphical range is long and it extends throughout the Mesozoic. *Brachyphyllum* in certain species may overlap the generic morphological features of *Pagiophyllum* (Harris, 1979).




Fig. 6. *Pagiophyllum* sites related to a reconstruction of Jurassic Pangea from literature (black dots) and Libyan site in (red dot) (Global Plate Tectonics, 2001).

Piazopteris branerii (White) Lorch, (Figs 7 c, d) Many fertile with very few fertile pinnae, age early Jurassic to Cretaceous, some of this Matoniaceae pinnulea probably belong to the *Weichaelia reticulata* (Stokes and Webb) Fontain which range in age from Middle Jurassic to Cenomanian, and it is commonly regarded as having been a plant of dry or salty habitats. This species is reported by Ash (1972) from the Lower Jurassic rocks, which was formerly thought to be of Triassic age near Ain Sukhana in the Gulf of Suez area of Egypt.

Hirmerella sp. (?) (Cone): There are several poorly preserved cones occurring in close association with the *Brachyphyllum* leafy shoots, and it is quite possible that some of them may belong to the *Hirmerella*, which is believed to be a member of the conifers that produced the peculiar pollen grains "Classopollis-type" in the Mesozoic time as suggested by Krassilov (1982). However, to make certain more materials need to be examined. If this proves to be true, then these are the first *Hirmerella* recorded from the African continent (C. Hill, personnel communication), thus they are of considerable interest. The *Hirmerella* group ranges in age from late Triassic "Rhaetian" to Early Cretaceous.

Otozamites sp. (leaf): This species occurs as small poorly preserved leaves. It ranges in age from Triassic to Early Cretaceous.

Samaropsis sp. (seed): They occur as seeds of uncertain classification, however, this species is known from Paleozoic, Mesozoic, and Cenozoic.




Fig. 7. SEM Latex replicas photos of a, b) *Brachyphyllum* sp. (stomata on the leaves arrowed; c, d) *Piazopteris branerii*.

4. Discussions

The *Pagiophyllum yafraiensis* sp. nov. is described and illustrated from the Khashm az Zarzur Formation for the first time. The described species is well preserved in a ferruginous clay of yellowish-brown color, soft, non-calcareous, non-swelling, with very rare quartz silt, few organic matters, and plant remains. The assigned age range of the reported associated plant assemblage from the previously published works is ranged from Jurassic to Early Cretaceous. However, none of the reported taxa (*Brachyphyllum* sp. *Piazopteris branerii*, *Hirmerella* sp., *Otozamites* sp., and *Samaropsis* sp.) are diagnostic of a specific age. The previous workers wrongly

determined some of these species and used them as strong stratigraphic markers, for example, *Piazopteris branerii* was incorrectly referred to as *Pecopteris* by Hammuda (1969, plate 2, figs. 13-15, p.69) (C. Hill personal communication).

The data on the flora in particular the co-occurrence of both *Pagiophyllum* and *Brachyphyllum* conifers as well as the sedimentological evidence suggest that warm and humid environments prevailed (El Zouki, 1980a). Furthermore, the abundance of conifers, particularly Podocarpacea (Chinnappa and Rajanikanth, 2017) with the taphocoenosis of the flora comprises local to regional elements derived from the riverbank, floodplain, back swamp, and valley settings (Fig. 8).

Burrollet (1963) assigned the Khashm az Zarzur to Late Jurassic (Callovian- Oxfordian) age based on the plant *Laccopteris saighanensis*. Hammuda (1969) reported from this rock unit *Pecopteris phillipsii*, *Neuropteris* sp., and *Zamites* sp. which point to the Jurassic age. In the Gulf of Suez area of Egypt Ash (1972) assigned the Early Jurassic age to rocks near Ain Sukhana due to the presence of *Piazopteris branerii*. A palynological investigation of a sample from a surface exposure in the vicinity of Yafran assigned the concerned Khashm az Zarzur Formation to Bathonian – Tithonian age (Hallett, 2002).




Fig. 8. Paleoeological reconstruction of Early Cretaceous flora from Pranhita-Godavari Basin (after Chinnappa and Rajanikanth, 2017).

5. Conclusions

- Six plant fossils taxa are reported from the Khashm az Zarzur Formation in the vicinity of Yafran, among which *Pagiophyllum yafraiensis* sp. nov. is described and illustrated herein.
- According to the sedimentological evidence and the associated floral assemblage, a warm with humid climate has been suggested.
- The collected fossil plants generally suggest a Jurassic to Lower Cretaceous age, but none of these species is diagnostic of a specific age. However, according to Burrollet (1963) and Hammuda (1969), the stratigraphic position of the exposed Khashm az Zarzur Formation "formerly known as Giosc Shale and Chameau Mort sandstone" which containing the *Pagiophyllum yafraiensis* sp. nov. is upper middle to lower early Jurassic (Callovian-Oxfordian) in age. However, based on palynological evidence the age was assigned to Bathonian-Tithonian (Hallett, 2002).

Acknowledgments

The British Museum is highly thankful for photographing and preparing the latex replica of the samples used in this paper. Dr. C. Hill of the British Museum (Natural History) of London is highly appreciated for his scientific help in the identification of the plant fossils.

Thanks are also extended to Dr. Milad Ben Rahuma of the Exploration Department, National Oil Corporation (NOC) for reviewing the stratigraphic chart.

References

- Ash, S.R., 1972. *Piazopteris branerii* from the Lower Jurassic, Egypt. Review of Palaeobotany and Palynology. 13 (2): 147-154.
- Antonovic, A. 1977. Sheet Mizdah (NH 33-1), Geological Map of Libya, Scale 1:250,000, Explanatory Booklet, Industrial Research Centre, Tripoli.
- Barr F.T., and Weegar, A.A., 1972. Stratigraphic Nomenclature of the Sirt Basin, Libya. Petrol. Explor. Soc. Libya, Tripoli, 179p.
- Banerjee, S., 1980. Stratigraphic lexicon of Libya. Ind. Res. Cent., Bull. 13:1-300.
- Burrollet, P.F., 1963. Field trip guidebook of the excursions to Jebel Nefusa (Mesozoic/Tertiary section in Tripolitania). Pet. Expl. Soc. Libya. 19p.
- Burrollet, P.F., 1977. Morphologie et pedologie d'une plaine couverte de la sable la Heffara Libyenne. Notes. Mèm. Comp. Fr. Petrole, 13:11-30.
- Cate, A., and Hammuda, O. S., 1969. Road log and map for Jebel Nefusa Field Trip. Geology of Libya Symposium. Department of Geology, University of Libya, 15p.
- Chinnappa, C., and Rajanikanth, A., 2017. Early Cretaceous flora from the Pranhita-Godavari Basin (east coast of India): taxonomic, taphonomic and paleoecological considerations. Acta Palaeobotanica 57 (1): 13-32.
- Christie, A. M., 1955. Geology of the Garian area, Tripolitania, Libya. U. N. Tech. Assistance Program Rept. TAA/LIB/2, 60p.
- Desio A., Ronchetti, R.C., Pozzi, R., Clerici, F. Invernizzi, G., Pisoni, C., and Vigano, F., 1963. Stratigraphic studies in the Tripolitanin Jebel (Libya). Rev. Ital. Paleontol. Stratigr. Mem 9, 126p.
- El-Hinnawy, M., and Cheshitey, G., 1975. Sheet Trabulus (NI 33-13), Geological Map of Libya. Scale 1:250,000, Explanatory Booklet, Industrial Research Centre, Tripoli.
- El Zouki, A. Y., 1976. Sedimentological and Stratigraphical aspects of the Chicla, Cabao and Chameau Mort formations (Upper Jurassic- Lower Cretaceous) of Jabal Nefusa, NW Libya. Ph.D. thesis, University of Bristol. 249p.
- El Zouki, A.Y., 1980a. Stratigraphy and Lithofacies of the Continental Clastic (Upper Jurassic and Lower Cretaceous) of Jabal Nafusah, NW Libya.. In: (M.J. Salem and M.T. Busrewil (eds) Second Symposium on the Geology of Libya. Academic Press, London. Vol. 3: 393-418.
- El Zouki, A.Y., 1980b. Depositional environment of the Chicla, Cabao, and Chameau Mort Formations as revealed by scanning electron microscopy. In: (M.J. Salem and M.T. Busrewil (eds) Second Symposium on the Geology of Libya. Academic Press, London. Vol. 3: 419-426.
- Fatmi, A. N., 1977. The Upper Triassic (Carnian) ammonite genus Mojsisovicsites from Jabal Gharuan, Libya. Libyan J. Sci. 7A:35-48.
- Fatmi, A.N., Sbata, A.M., and Eliagoubi, B.A. 1978. Guide to the Mesozoic stratigraphy of Jabal Nefusa, Libyan Jamahiriya. Publication No. 7, Arab Development Inst, Tripoli. 35p.
- Fatmi, A.N., Eliagoubi, B.A., and Hammuda, O.S. 1980. Stratigraphic nomenclature of the pre-Upper Cretaceous Mesozoic rocks of Jabal Nafusah, NW. Libya. Second Symposium on the Geology of Libya, vol. 1 (eds. M.J. Salem and M.T. Busrewil), Academic Press, London, p. 57-66.
- Global Plate Tectonics (2001) I Geology B Plate Tectonics 2 Midocean Ridge, Plates/UTIG (2001), from <https://slidetodoc.com/i-geology-b-plate-tectonics-2-midocean-ridge/>.

- Hallett, D., 2002. Petroleum Geology of Libya. Elsevier B. V. Amsterdam, The Netherlands. 503p.
- Hammuda, O.S., 1969. Jurassic and Lower Cretaceous rocks of central Jabal Nefusa, NW Libya. Petroleum Exploration Society of Libya. 74p.
- Harris, T. M., 1979. The Yorkshire Jurassic Flora. V. Voniferales, Br. Mus. (Nat. Hist.), 803, 166p.
- Krassilov, V.A., 1982. On the ovuliferous organ of Hirmeriella. *Phyta* 1 : 141 – 144.
- Mann, K. 1975. Sheet Al Khums (NI 33-14), Geological Map of Libya, scale 1:250,000, Explanatory Booklet, Industrial Research Centre, Tripoli.
- Novović, T. 1977. Sheet Nalut (NH 32-4). Geological Map of Libya, scale 1:250,000, Explanatory Booklet. Industrial Research Centre, Tripoli.
- Shiref, Y.M., and Salaj, J., 2007. Upper Jurassic—Lower Paleogene lithostratigraphy and facies development in the Al Hamadah al Hamra area (Libya). *Geologica Carpathica*, 58 (1), 3-18.
- Smetana, R. 1975. Sheet Ra's Jdeir (NI 32-16), Geological Map of Libya, scale 1:250,000, Explanatory Booklet, Industrial Research Centre, Tripoli.
- Stockey, R.A., 1982. The Araucariaceae: An evolutionary perspective. *Review of Palaeobotany and Palynology*, 37: 133-154.
- Taylor, T.N., Taylor, E.L., and Krings, M., 2009. The biology and evolution of fossil plants, 2nd ed. Academic Press, Amsterdam, Netherlands.
- Živanović, M. 1977. Sheet Bani Walid (NH 33-2), Geological Map of Libya, Scale 1:250,000, Explanatory Booklet, Industrial Research Centre, Tripoli.
- <https://en.wikipedia.org/wiki/pagiophyllum>